

 **Kirsty™**  
(insulin aspart-xjhz) injection  
100 units/mL (U-100)

**1<sup>ST</sup>** Interchangeable  
biosimilar to NovoLog®<sup>1</sup>



## Satisfy Your Mealtime Insulin Needs<sup>2</sup>

### **IMPORTANT SAFETY INFORMATION**

**Do not share your KIRSTY Prefilled Pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.**

**Do not take KIRSTY if you:**

- are having an episode of low blood sugar (hypoglycemia).
- have an allergy to insulin aspart products or any of the ingredients in KIRSTY.

**Please see additional Important Safety Information on pages 10-11 and accompanying Full Prescribing Information.**

# KIRSTY CAN.



Give Mealtime  
Insulin Control<sup>2</sup>



Deliver Clinical  
Confidence<sup>2</sup>



Provide Savings  
& Support

With Kirsty™, a biosimilar option that works like NovoLog (insulin aspart), more is possible in managing diabetes.<sup>2</sup>

2

 **Kirsty™**  
(insulin aspart-xjhz) injection  
100 units/mL (U-100)



You can also visit us  
at [Kirsty.com](https://www.kirsty.com) to learn more.

### **IMPORTANT SAFETY INFORMATION (continued)**


Before taking KIRSTY, tell your healthcare provider about all your medical conditions, including if you are:

- pregnant, planning to become pregnant, or are breastfeeding.
- taking new prescription or over-the-counter medicines, vitamins, or herbal supplements.

Please see additional Important Safety Information on pages 10-11 and accompanying Full Prescribing Information.

# SERVING SIMILARITY YOU CAN TRUST.

KIRSTY is a rapid-acting insulin biosimilar that's interchangeable with NovoLog from a trusted name in insulin, Biocon Biologics.<sup>1,2</sup>



Similar Safety<sup>2-4</sup>



Similar Effectiveness<sup>2-4</sup>

 **Kirsty™**  
(insulin aspart-xjhz) injection  
100 units/mL (U-100)

## What is a biosimilar?

A **biosimilar** is a nearly identical copy of an existing medicine. It works in the same way and is as safe and effective, so you can be sure to receive the same care as with the original approved medicine.<sup>3</sup>

## What is an *interchangeable* biosimilar?

An **interchangeable biosimilar** meets the same high standards of biosimilarity and can be safely switched from the original approved medicine to a biosimilar by a pharmacist without doctor approval—helping more people get the medicine they need.\*<sup>3</sup>

\*Subject to state pharmacy laws.



Similar Prefilled Pen<sup>2-4</sup>

## IMPORTANT SAFETY INFORMATION (continued)

**Before you start taking KIRSTY, talk to your healthcare provider about low blood sugar and how to manage it.**

**KIRSTY starts acting fast.** You should eat a meal within 5 to 10 minutes after you take your dose of KIRSTY.


**Check your blood sugar levels.**

**Do not reuse or share your needles with other people.**

**Please see additional Important Safety Information on pages 10-11 and accompanying Full Prescribing Information.**

# HELPING YOU SAFELY INJECT KIRSTY.

Whether it's your first time or you need a little refresher, our instructional video will help you see how simple using the KIRSTY prefilled pen can be.



We'll walk you through the injection steps so that each dose of KIRSTY feels more like a normal part of your daily routine.



Scan or visit  
[www.Kirsty.com/#how-to-inject](http://www.Kirsty.com/#how-to-inject)  
to watch our quick video tutorial.

 **Kirsty™**  
(insulin aspart-xjhz) injection  
100 units/mL (U-100)

KIRSTY is available in 2 forms<sup>2</sup>



## KIRSTY Prefilled Pen

Visit [Kirsty.com](http://Kirsty.com) to download the  
Instructions for Use



## KIRSTY 10 mL Vial

Visit [Kirsty.com](http://Kirsty.com) to download the  
Instructions for Use

Dosing should be the same when changing from NovoLog to KIRSTY.<sup>2,4</sup> Be sure to check with your healthcare provider to confirm your dose.

### **IMPORTANT SAFETY INFORMATION (continued)**

KIRSTY can be injected under the skin (subcutaneously) of your stomach area (abdomen), buttocks, upper legs (thighs) or upper arms, or by continuous infusion under the skin (subcutaneously) through an insulin pump into an area of your body recommended in the instructions that come with your insulin pump.

**Please see additional Important Safety Information on pages 10-11 and accompanying Full Prescribing Information.**

# DELIVERING CARE

Commercially insured for KIRSTY?  
Our support offerings can help you more easily access your medicine when you need it.

## Trial Voucher Program\*

The Trial Voucher Program offers eligible patients a voucher for a free trial of 5 pens or 2 vials of insulin.

- The voucher must be requested by a healthcare professional
- A valid prescription must accompany the voucher

\*Restrictions apply. One voucher redemption per patient. Please see full terms and conditions at [www.Kirsty.com/trialvoucherprogram](http://www.Kirsty.com/trialvoucherprogram).

# WITHIN YOUR REACH.

## Savings Card Program\*

- Biocon Biologics offers copay support for eligible patients to **pay as little as \$35** in out-of-pocket costs

\*Eligibility restrictions apply. Not valid for uninsured patients who are covered by a state- or federally funded healthcare program. Please see full terms and conditions at [www.Kirsty.com/copay](http://www.Kirsty.com/copay).



For questions regarding our support services, call 1-800-657-7613  
Monday-Friday 8 AM-8 PM ET.

Find out if you're eligible for copay assistance and get enrolled by visiting  
[www.Kirsty.com/copayenroll](http://www.Kirsty.com/copayenroll).

## IMPORTANT SAFETY INFORMATION

### What is KIRSTY (insulin aspart-xjhz) injection 100 Units/mL?

- KIRSTY is a prescription man-made insulin that is used to control high blood sugar in adults and children with diabetes mellitus.

## IMPORTANT SAFETY INFORMATION

**Do not share your KIRSTY Prefilled Pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.**

### Do not take KIRSTY if you:

- are having an episode of low blood sugar (hypoglycemia).
- have an allergy to insulin aspart products or any of the ingredients in KIRSTY.

**Before taking KIRSTY, tell your healthcare provider about all your medical conditions, including if you are:**

- pregnant, planning to become pregnant, or are breastfeeding.
- taking new prescription or over-the-counter medicines, vitamins, or herbal supplements.

**Before you start taking KIRSTY, talk to your healthcare provider about low blood sugar and how to manage it.**

**KIRSTY starts acting fast.** You should eat a meal within 5 to 10 minutes after you take your dose of KIRSTY.

**Check your blood sugar levels.**

**Do not reuse or share your needles with other people.**

KIRSTY can be injected under the skin (subcutaneously) of your stomach area (abdomen), buttocks, upper legs (thighs) or upper arms, or by continuous infusion under the skin (subcutaneously) through an insulin pump into an area of your body recommended in the instructions that come with your insulin pump.

**Change (rotate) your injection sites within the area you choose with each dose** to reduce your risk of getting lipodystrophy (pits in skin or thickened skin) and localized cutaneous amyloidosis (skin with lumps) at the injection sites. **Do not** use in the exact same spot for each injection, inject where the skin has pits, is thickened or has lumps. **Do not** inject where the skin is tender, bruised, scaly, hard, into scars or damaged skin.

While taking KIRSTY do not drive or operate heavy machinery, until you know how KIRSTY affects you. You should not drink alcohol or use prescription or over-the-counter medicines that contain alcohol.

KIRSTY may cause serious side effects that can lead to death, including, low blood sugar (hypoglycemia). Signs and symptoms that may indicate low blood sugar include dizziness or light-headedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion, shakiness, headache or fast heartbeat.

Your insulin dose may need to change because of change in level of physical activity or exercise, increased stress, change in diet, weight gain or loss or illness.

Other common side effects of KIRSTY may include low potassium in your blood (hypokalemia), reactions at the injection site, itching, rash, serious allergic reactions (whole body reactions), skin thickening or pits at the injection site (lipodystrophy), weight gain, and swelling of your hands and feet.

**Get emergency medical help if you have** trouble breathing, shortness of breath, fast heartbeat, swelling of your face, tongue, or throat, sweating, extreme drowsiness, dizziness, confusion.

**These are not all the possible side effects of KIRSTY. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**References:** **1.** Jeremias S. FDA approves Kirsty as first interchangeable insulin aspart biosimilar. The Center for Biosimilars. July 15, 2025. Accessed July 31, 2025. <https://www.centerforbiosimilars.com/view/fda-approves-kirsty-as-first-interchangeable-insulin-aspart-biosimilar> **2.** KIRSTY. Prescribing information. Biocon Biologics Inc; 2025. **3.** US Food and Drug Administration. Biosimilars: review and approval. Last updated December 13, 2022. Accessed July 31, 2025. <https://www.fda.gov/drugs/biosimilars/review-and-approval> **4.** NovoLog. Prescribing Information. Novo Nordisk Inc; 2023.





 **Kirsty™**  
(insulin aspart-xjhz) injection  
100 units/mL (U-100)

**KIRSTY** can because

**BIOCON  
BIOLOGICS  
CAN.**

Learn more about us at [BioconBiologicsUS.com](https://BioconBiologicsUS.com).

KIRSTY and the Kirsty Logo are trademarks of Biocon Biologics Limited.  
BIOCON BIOLOGICS and the Biocon Biologics Logo are registered  
trademarks of Biocon Biologics Limited.

NovoLog is a registered trademark of Novo Nordisk A/S.

© 2025 Biocon Biologics Inc. All Rights Reserved. ASP-2025-00014-US

 **Biocon Biologics®**